



The use of a plant enzyme provides fast, sensitive and reproducible LC-MS/MS method for folate analyses in food

Ložnjak, Petra; Jakobsen, Jette; Bysted, Anette; Diaz de la Garza, Rocio; Rychlik, Michael

Published in:
5th International Vitamin Conference - book of abstract

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Ložnjak, P., Jakobsen, J., Bysted, A., Diaz de la Garza, R., & Rychlik, M. (2018). The use of a plant enzyme provides fast, sensitive and reproducible LC-MS/MS method for folate analyses in food. In *5th International Vitamin Conference - book of abstract* (pp. 58-58). [29].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract no.: 29

Title: The use of a plant enzyme provides a fast, sensitive and reproducible LC-MS/MS method for folate analyses in food

Petra Loznjak^{1,2}, Jette Jakobsen^{1,2}, Anette Bysted^{1,2}, Rocio Isabel Diaz de la Garza^{3,4} and Michael Rychlik^{5,6}

¹ *National Food Institute*, ² *Technical University of Denmark*, ³ *FEMSA Biotechnology Center*, ⁴ *Institute of Technology and Higher Education*, ⁵ *Chair of Analytical Food Chemistry*, ⁶ *Technical University of Munich*

Folate analyses in food are one of the most challenging due to the structural variability, low stability and low presence of folates, but also because of the complexity of food matrices. The use of protease, amylase and deconjugase of animal origin, known as a tri-enzyme extraction, has been considered as a necessity in order to provide the complete determination of folates in food. However, there are discrepancies in the data for folate content between methods due to the lack of a proper deconjugase enzyme which would enable fast and reproducible folate deglutamylation.

The newly developed method included single-enzymatic extraction step in which an enzyme of a plant origin (*Arabidopsis thaliana*) has been used. Seven folate vitamers (tetrahydrofolate, 5,10-methenyl-tetrahydrofolate, 10-formyl-folic acid, 5-formyl-tetrahydrofolate, folic acid, 5-methyl-tetrahydrofolate and 10-methyl-folic acid) were quantified using the ¹³C₅-labelled internal standards by LC-MS/MS method. Deconjugase of a plant origin provided sufficient folate deglutamylation within 1 hour, showing the activity >95%. Consequently, it enabled performance of the whole analysis within 1 working day, omitting commonly used overnight incubation. The method was validated to assess precision ($\pm 10\%$ RSD) and trueness (80-120%). Additionally, 89 samples of dairy products, fruit, vegetables, legumes, cereals, offal and meat were analyzed with this new method, another LC-MS/MS method using rat serum deconjugase, and the traditional microbiological assay.

Furthermore, the mobile phases, column and the MS-ion source used in this method, may be used for determination of vitamins D and K, which provides an easy shift in the analyses of these vitamins in the lab.